

An unusual cause of dyspepsia due to a retained epicardial pacing wire

Gökhan Aksakal¹, Emre Kubat², Mehmet Ali Erdoğan¹, Celal Secuk Unal²

(1) Karabuk Training and Research Hospital, Department of Gastroenterology ; (2) Karabuk Training and Research Hospital, Department of Cardiovascular Surgery.

To the Editor,

Temporary epicardial pacing wires (TEPWs) are frequently inserted in cardiac surgery procedures. Although overall morbidity related to TEPWs is low, complications related to the insertion, utilization, retrieval, and migration of these wires have been reported in the literature (1). In this report, we describe an unusual case of gastric migration of a TEPW, which eventually caused dyspeptic symptoms.

A 74-year-old man was admitted to our gastroenterology clinic for evaluation of worsening dyspeptic symptoms. Over the previous four months, he had suffered from continuous epigastric burning and pain. The patient had undergone coronary bypass grafting one year prior to the admission. On upper gastrointestinal endoscopy, a blue wire was identified that had penetrated the distal anterior wall of the gastric corpus (Figure 1). Retrieval of the foreign body was attempted but failed and the endoscopic procedure was terminated. For further evaluation, thoracic and abdominal computed tomography scans were performed. With imaging, the location and path of the wire were confirmed, from the right ventricle, passing through the sternocostal triangle, entering the abdomen, and penetrating the gastric wall (Figure 2). Consultation with cardiovascular surgery confirmed this wire to be a TEPW, which was cut at the surface of the skin after the bypass procedure.

Dysrhythmia frequently occurs in the postoperative period of cardiac surgery. Temporary pacing is a simple and efficient method for treating dysrhythmia, which causes hemodynamic instability (2). When used, TEPWs are commonly removed on the fourth postoperative day with gentle transcutaneous traction. Rarely, the wires are cut at the surface of the skin, with the risk of retraction into the subcutaneous tissues due to the residual traction on the wire and/or increased coagulation rates. Thoracic complications during retrieval are the most common catastrophic complications of TEPWs (2). Retained wires also carry the potential risk for infection. However, major abdominal complications related to retained TEPWs have rarely been reported in the literature. Acute colonic injury, abdominal aortic migration, and colonic migration have been reported as TEPW-related abdominal complications (3-5). To the best of our knowledge, our case is the first report of gastric wall injury secondary to the migration of TEPW into the gastric cavity. We believe that the retained wire migrated in the abdominal cavity and penetrated the

gastric wall, leading to chronic dyspeptic symptoms, 10 months after cardiac surgery.

In conclusion, we believe that these complications are mostly associated with improper placement of the TEPWs. Furthermore, the wire should not be passed through the intraabdominal cavity to prevent this




Fig. 1. — Upper gastrointestinal endoscopy showing a penetrated wire in the anterior wall of corpus.

Correspondence to : Gökhan Aksakal, Karabuk Training and Research Hospital, Department of Gastroenterology, Alparslan street, Karabuk, Turkey.
E-mail : gkhnsk1@gmail.com.

Submission date : 06/05/2016

Acceptance date : 31/05/2016




Fig. 2. — Computed tomography images showing a wire located at the right ventricle, passing through the sternocostal triangle, entering the abdomen, and penetrating the gastric wall (arrowhead).

complication, which could be a cause of morbidity or mortality in patients following cardiac surgery.

References

1. ABU-OMAR Y., GUERRIERI-WOLF L., TAGGART D.P. Indications and positioning of temporary pacing wires. *Multimed. Manual Cardiothorac. Surg.*, 2006, 2006.0512, mmcts.2005.001248.
2. DIMITRAKAKIS G., PODILA S.R., MEHTA D. Re : Temporary epicardial pacing wire removal: not an innocuous procedure. *Interact. Cardiovasc. Thorac. Surg.*, 2010, **11** : 855-856.
3. SALAMI M.A., COLEMAN R.J., BUCHAN K.G. Colonic injury from temporary epicardial pacing wires. *Ann. Thorac. Surg.*, 2012, **93** : 1309-1311.
4. MUKAIHARA K., YOTSUMOTO G., MATSUMOTO K., IMOTO Y. Migration of a retained temporary epicardial pacing wire into an abdominal aortic aneurysm. *Eur. J. Cardiothorac. Surg.*, 2015, **48** : 169-170.
5. GONZALES S., WHITE H., ECHAVARRIA J. Transcolonic migration of retained epicardial pacing wires. *Case Rep. Radiol.*, 2015 : 416587.