

## A rare cause of colic polyp in a patient with ischemic colitis

E. Curraj<sup>1</sup>, R. Ntounda<sup>2</sup>, B. Van Houte<sup>2</sup>, G. Delporte<sup>3</sup>, D. Lacreman<sup>3</sup>

(1) Department of Internal Medicine, Cliniques Universitaires Saint-Luc, Bruxelles, Belgium ; (2) Department of Gastroenterology, Centre Hospitalier Régional du Val de Sambre, Auvelais, Belgium ; (3) Department of Pathology, Centre Hospitalier Universitaire Liege, Liège, Belgium.

### Quiz

A 50-year-old woman was admitted for a rectal bleeding after an episode of defecation 2 days prior to her admission. The patient had also suffered from lower abdominal pain for 3 days, but did not complain about any other pain.

The patient had no personal or familial medical history.

The clinical examination only revealed a sensitive abdomen in the left inferior quadrant of the abdomen without defence or rebound. The vital parameters were normal.

The blood sample showed the following: haemoglobin 15.2 g/dL (12-16), MCV 105 (85-95), leucocytes 11800/mm<sup>3</sup> (4200-9000) among which 8470/mm<sup>3</sup> neutrophils (1900-7000), platelets 225000/mm<sup>3</sup> (150000-400000) and CRP 171.5 mg/dL (< 12). We noticed that the haemoglobin had decreased from 15.2 g/dL to 11.6 g/dL between the emergency and our gastroenterological department.

The abdominal CT contrast showed a sub mucosal thickening of the left colic, sigmoid and rectal walls, and a para-colic infiltration.

The sigmoidoscopy performed 4 days after admission revealed an ischemic colitis and a submucosal polyp in the middle sigmoid.

Eight weeks later the patient came back for a total colonoscopy. She was asymptomatic. The exams enabled the complete mucosal en bloc resection of the submucosal polyp located 70 cm away from the anal margin. It was a white oval and mobile polyp of 2cm in size and of hard consistency.

Based on the aforementioned description, what is the nature of the polyp?

### Answer

The pathological analysis with immunohistochemistry showed a mesenchymal benign tumour of 6mm with a diffuse marking inside the lesion with Protein S100, a heterogeneous marking with Synaptophysin and a cytoplasmic granular marking with PAS + Diastase.

We reached the diagnosis of ganglioneuroma, a peripheral and benign nerve tumour which is part of the neuroblastic tumours. These tumours are mostly retroperitoneal (45-50%) and mediastinal (40%). The excision with complete resection is essential because of the tumour's malignant capacity.




Figure 1. — White oval polyp of 2cm size at the control colonoscopy (8 weeks after the sigmoidoscopy).




Figure 2. — Pathological analysis with immunohistochemistry (1) Histology; (2) Protein S100; (3) Synaptophysin.

### Conflict of interest

The authors have no conflict of interest.

Correspondence to: Edwin Curraj, Rue Jacques Ballings, 19, 1140 Evere, Belgium.  
E-mail: edwincurraj@gmail.com

Submission date: 21/12/2018  
Acceptance date: 02/04/2019

*Acta Gastro-Enterologica Belgica*, Vol. LXXXII, July-September 2019